



PROVEN



CITRATE-FREE



PFIZER SERVICES &
SUPPORT

Your guide to treatment with ABRILADA. Let's go.

Injection: 40 mg/0.8 mL, 20 mg/0.4 mL

Pfizer's FDA-approved interchangeable biosimilar to Humira[®] (adalimumab)*

Not actual patient.

*ABRILADA is interchangeable for the indications of use, dosage forms, strengths and routes of administration described in [Prescribing Information](#).

SELECTED SAFETY INFORMATION

What is the most important information I should know about ABRILADA?

You should discuss the potential benefits and risks of ABRILADA with your doctor. ABRILADA is a tumor necrosis factor (TNF) blocker medicine that can lower the ability of your immune system to fight infections. You should not start taking ABRILADA if you have any kind of infection unless your doctor says it is okay.

- **Serious infections have happened in people taking adalimumab products. These serious infections include tuberculosis (TB) and infections caused by viruses, fungi, or bacteria that have spread throughout the body. Some people have died from these infections.** Your doctor should test you for TB before starting ABRILADA, and check you closely for signs and symptoms of TB during treatment with ABRILADA, even if your TB test was negative. If your doctor feels you are at risk, you may be treated with medicine for TB.
- **Cancer.** For children and adults taking TNF blockers, including ABRILADA, the chance of getting lymphoma or other cancers may increase. There have been cases of unusual cancers in children, teenagers, and young adults using TNF blockers. Some people have developed a rare type of cancer called hepatosplenic T-cell lymphoma. This type of cancer often results in death. If using TNF blockers including ABRILADA, your chance of getting two types of skin cancer (basal cell and squamous cell) may increase. These types are generally not life-threatening if treated; tell your doctor if you have a bump or open sore that doesn't heal.

Please see Selected Safety Information throughout, Indications on page 3, and [Full Prescribing Information, including BOXED WARNING, Medication Guide, and Instructions for Use](#), available at [ABRILADA.com](#).



Not actual patients.

What is ABRILADA?

ABRILADA is a biologic treatment that is approved for certain inflammatory conditions (listed on the right). ABRILADA is a biosimilar—a highly similar version of Humira® (adalimumab), a biologic medicine that has been available for some time.

A biosimilar is approved by the US Food and Drug Administration (FDA) only after it has been found to have no clinically meaningful differences from the original biologic medicine.

Whether you are starting on ABRILADA or switching from Humira, this brochure can help you understand your treatment.

Be sure to review the benefits and risks of ABRILADA, discuss them with your doctor, and read the [Medication Guide](#) and [Instructions for Use](#) before starting treatment.

In this brochure, you'll also find information on possible side effects that are written in the form of questions. Talk to your doctor about these and any other questions you may have about ABRILADA.

To help you get started on your ABRILADA treatment journey, this brochure also talks about resources and potential financial support from Pfizer.

SELECTED SAFETY INFORMATION

What should I tell my doctor BEFORE starting ABRILADA?

Tell your doctor about all of your health conditions, including if you:

- Have an infection, are being treated for infection, or have symptoms of an infection
- Get a lot of infections or infections that keep coming back
- Have diabetes
- Have TB or have been in close contact with someone with TB, or were born in, lived in, or traveled where there is more risk for getting TB

What conditions is ABRILADA approved for?

ABRILADA is a medicine called a tumor necrosis factor (TNF) blocker that is used to reduce the signs and symptoms of:

Moderate to severe rheumatoid arthritis (RA) in adults. ABRILADA can be used alone, with methotrexate, or with certain other medicines.

Moderate to severe polyarticular juvenile idiopathic arthritis (JIA) in children 2 years and older. ABRILADA can be used alone or with methotrexate.

Psoriatic arthritis (PsA) in adults. ABRILADA can be used alone or with certain other medicines.

Ankylosing spondylitis (AS) in adults.

Moderate to severe hidradenitis suppurativa (HS) in adults.

ABRILADA is also used to treat:

Moderate to severe Crohn's disease (CD) in adults and children 6 years of age and older.

Moderate to severe ulcerative colitis (UC) in adults.

It is not known if adalimumab products are effective in people who stopped responding to or could not tolerate TNF-blocker medicines.

Moderate to severe chronic (lasting a long time) plaque psoriasis (Ps) in adults who have the condition in many areas of their body, and who may benefit from taking injections or pills (systemic therapy), or phototherapy (treatment using ultraviolet light, alone or with pills).

Uveitis (UV) in adults. Non-infectious intermediate (middle part of the eye), **posterior** (back of the eye), and **panuveitis** (all parts of the eye).

*Please see Selected Safety Information throughout, Indications above, and Full Prescribing Information, including **BOXED WARNING**, Medication Guide, and Instructions for Use, available at ABRILADA.com.*

How does ABRILADA work?

ABRILADA blocks the action of a protein in the body called tumor necrosis factor-alpha (TNF-alpha). TNF-alpha is made by your body's immune system. People with certain diseases have too much TNF-alpha, which can cause the immune system to attack normal healthy parts of the body. ABRILADA works by blocking TNF-alpha, which in excess amounts can cause inflammation.

How is ABRILADA administered?

ABRILADA is given by an injection under the skin. Your healthcare professional (HCP) will tell you how often to take an injection of ABRILADA.

Do not inject ABRILADA more often than your HCP has instructed.

How will I get my ABRILADA prescription?

Your ABRILADA prescription will most likely be filled by a specialty pharmacy.

A specialty pharmacy provides medications used to treat complex, chronic health problems.

These medicines may also need special handling by the pharmacist and you (special storage or detailed treatment planning, for example).

Your insurance provider will select a specialty pharmacy for you. The specialty pharmacy will:


- Prepare your ABRILADA medicine
- Contact you when it is ready
- Ship it directly to you

SELECTED SAFETY INFORMATION

What should I tell my doctor BEFORE starting ABRILADA? (continued)

Tell your doctor about all of your health conditions, including if you: (continued)

- Live or have lived in an area (such as the Ohio and Mississippi River valleys) where there is an increased risk for getting certain kinds of fungal infections, such as histoplasmosis, coccidioidomycosis, or blastomycosis. These infections may happen or become more severe if you use ABRILADA. Ask your doctor if you are unsure if you have lived in these areas
- Have or have had hepatitis B
- Are scheduled for major surgery
- Have or have had cancer



Not actual patient.

See the full [Medication Guide](#) and [Instructions for Use](#) available at ABRILADA.com for complete instructions on how to properly prepare and inject ABRILADA

SELECTED SAFETY INFORMATION

What should I tell my doctor BEFORE starting ABRILADA? (continued)

Tell your doctor about all of your health conditions, including if you: (continued)

- Have numbness or tingling or a nervous system disease such as multiple sclerosis or Guillain-Barré syndrome
- Have or had heart failure
- Have recently received or are scheduled to receive a vaccine. ABRILADA patients may receive vaccines, except for live vaccines. Children should be brought up to date on all vaccines before starting ABRILADA
- Are allergic to any ABRILADA ingredients

Please see Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including BOXED WARNING, Medication Guide, and Instructions for Use, available at ABRILADA.com.

INJECTION
Abrilada[™]
adalimumab-afzb

Not actual patients.

If you are new to biologic therapy

ABRILADA is a biologic treatment that may help with a variety of conditions (see page 3 for the conditions ABRILADA is approved for). Your doctor will decide if treatment with ABRILADA is appropriate for you.

During your visit to your HCP's office, you may learn more about:

- ✓ How ABRILADA works and possible side effects
- ✓ How often to inject ABRILADA
- ✓ How Pfizer enCompass[®] can help assess your insurance coverage for ABRILADA and answer additional coverage-related questions you may have

Please refer to the *Some questions to ask your doctor* section on pages 10-13 for more information about treatment with ABRILADA—and be sure to review the benefits and risks of ABRILADA with your doctor (see safety information on pages 14 and 15).

SELECTED SAFETY INFORMATION

What should I tell my doctor BEFORE starting ABRILADA? (continued)

Tell your doctor about all of your health conditions, including if you: (continued)

- Are pregnant, planning to become pregnant, breastfeeding, or planning to breastfeed
- Have a baby and you were using ABRILADA during your pregnancy. Tell your baby's doctor before your baby receives any vaccines

Also tell your doctor about all the medicines you take. You should not take ABRILADA with ORENCIA[®] (abatcept), KINERET[®] (anakinra), REMICADE[®] (infliximab), ENBREL[®] (etanercept), CIMZIA[®] (certolizumab pegol), or SIMPONI[®] (golimumab). Tell your doctor if you have ever used RITUXAN[®] (rituximab), IMURAN[®] (azathioprine), or PURINETHOL[®] (mercaptapurine, 6-MP).

How do I take ABRILADA?

ABRILADA provides patients with 2 options for self-administration: the prefilled, single-dose pen and the prefilled, single-dose syringe.

Before your injection, your doctor will decide the right dose of ABRILADA and how often you should receive it. Do not inject ABRILADA more often than you were prescribed.

Do not try to inject ABRILADA until you are shown how to give the injections and have read and understand the [Instructions for Use](#). If your healthcare professional decides that you or a caregiver may be able to give your injections of ABRILADA at home, you or your caregiver should receive education on how to do so.

ABRILADA is designed with your needs in mind

Citrate-free

- ABRILADA is citrate-free, which may reduce injection site pain

Made without natural rubber latex

- This may reduce the likelihood of having an allergic reaction

Out-of-refrigerator stability

- ABRILADA must be refrigerated at 36°F to 46°F. If needed, for example when traveling, ABRILADA can stay outside the refrigerator for up to 30 days (at a maximum of 86°F), out of direct sunlight



Download the Travel Guidance Brochure from ABRILADA.com for helpful tips that may make traveling with ABRILADA easier.

SELECTED SAFETY INFORMATION

What should I watch for AFTER starting ABRILADA?

ABRILADA can cause serious side effects, including:

- **Serious infections.** These include TB and infections caused by viruses, fungi, or bacteria. Symptoms related to TB include a cough, low-grade fever, weight loss, or loss of body fat and muscle.

Call your doctor or get medical care right away if you develop any of the above symptoms.

*Please see Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including **BOXED WARNING, Medication Guide, and Instructions for Use**, available at ABRILADA.com.*

INJECTION
Abrilada[™]
 adalimumab-afzb



Not actual patients.

If you are switching to ABRILADA from Humira

If you are switching to ABRILADA from Humira® (adalimumab), you can expect it to provide similar treatment outcomes to what you had with Humira. Individual results may vary.

With ABRILADA, you can expect a similar treatment experience as Humira

- ✓ Works the same way as Humira
- ✓ Dosed the same way as Humira
- ✓ Is approved for all eligible indications of Humira*

*ABRILADA is approved for all indications of Humira except for pediatric UC, adolescent HS, and pediatric UV.

SELECTED SAFETY INFORMATION

What should I watch for AFTER starting ABRILADA? (continued)

ABRILADA can cause serious side effects, including: (continued)

- **Hepatitis B infection in carriers of the virus.** Symptoms include muscle aches, feeling very tired, dark urine, skin or eyes that look yellow, little or no appetite, vomiting, clay-colored bowel movements, fever, chills, stomach discomfort, and skin rash.
- **Allergic reactions.** Symptoms of a serious allergic reaction include hives, trouble breathing, and swelling of your face, eyes, lips, or mouth.
- **Nervous system problems.** Signs and symptoms include numbness or tingling, problems with your vision, weakness in your arms or legs, and dizziness.

Call your doctor or get medical care right away if you develop any of the above symptoms.

Why should I switch to ABRILADA?

Your doctor may have decided that ABRILADA is appropriate to treat your condition. Because Humira may be working well, it's important to know that you can expect similar safety and efficacy with ABRILADA as you had with Humira.

Will the dose I receive change with ABRILADA?

Your doctor will determine the appropriate dose of ABRILADA for you and how often you should receive it. This will be based on your medical condition.

Please refer to the *Some questions to ask your doctor* section on pages 10-13 for more information about your treatment with ABRILADA—and be sure to review the benefits and risks of ABRILADA with your doctor.

Pfizer enCompass® can help assess your insurance coverage for ABRILADA and answer additional coverage-related questions you may have

SELECTED SAFETY INFORMATION


What should I watch for AFTER starting ABRILADA? (continued)

ABRILADA can cause serious side effects, including: (continued)

- **Blood problems** (decreased blood cells that help fight infections or stop bleeding). Symptoms include a fever that does not go away, bruising or bleeding very easily, or looking very pale.
- **Heart failure** (new or worsening). Symptoms include shortness of breath, swelling of your ankles or feet, and sudden weight gain.
- **Immune reactions including a lupus-like syndrome.** Symptoms include chest discomfort or pain that does not go away, shortness of breath, joint pain, or rash on your cheeks or arms that gets worse in the sun.
- **Liver problems.** Symptoms include feeling very tired, skin or eyes that look yellow, poor appetite or vomiting, and pain on the right side of your stomach (abdomen). These problems can lead to liver failure and death.
- **Psoriasis** (new or worsening). Symptoms include red scaly patches or raised bumps that are filled with pus.

Call your doctor or get medical care right away if you develop any of the above symptoms.

Please see *Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including BOXED WARNING, Medication Guide, and Instructions for Use, available at ABRILADA.com.*



Not actual patient or healthcare professional.

Some questions to ask your doctor

Whether you are new to adalimumab or switching from Humira[®] (adalimumab), it is important to review the safety information with your HCP, including potential side effects and medications to avoid during treatment. Be sure to review the benefits and risks of ABRILADA with your doctor and read the [Medication Guide](#) before starting treatment.

Are there any side effects I should know about before starting treatment with ABRILADA?

ABRILADA can cause serious side effects, including:

• Cancer

- For children and adults taking TNF blockers, including ABRILADA, the chances of getting cancer may increase
- There have been cases of unusual cancers in children, teenagers, and young adults using TNF blockers
- People with RA, especially more serious RA, may have a higher chance for getting a kind of cancer called lymphoma
- If you use TNF blockers, including ABRILADA, your chance of getting 2 types of skin cancer may increase (basal cell cancer and squamous cell cancer of the skin). These types of cancer are generally not life-threatening if treated. Tell your HCP if you have a bump or open sore that does not heal

— Some people receiving TNF blockers, including ABRILADA, developed a rare type of cancer called hepatosplenic T-cell lymphoma. This type of cancer often results in death. Most of these people were male teenagers or young men. Also, most people were being treated for Crohn's disease or ulcerative colitis with another medicine called Imuran[®] (azathioprine) or Purinethol[®] (6-mercaptopurine, 6-MP)

• Serious infections

Your HCP will examine you for tuberculosis (TB) and perform a test to see if you have TB. If your HCP feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with ABRILADA and during treatment with ABRILADA.

Even if your TB test is negative your HCP should carefully monitor you for TB infections while you are taking ABRILADA. People who had a negative TB skin test before receiving adalimumab products have developed active TB.

Tell your HCP if you have any of the following symptoms while taking or after taking ABRILADA:

- Cough that does not go away
- Low-grade fever
- Weight loss
- Loss of body fat and muscle (wasting)

• Hepatitis B infection in people who carry the virus in their blood

Tell your HCP if you have any of the following symptoms of a possible hepatitis B infection:

- Muscle aches
- Feel very tired
- Dark urine
- Skin or eyes look yellow
- Little or no appetite
- Vomiting
- Clay-colored bowel movements
- Fever
- Chills
- Stomach discomfort
- Skin rash

• Allergic reactions

Allergic reactions can happen in people who use ABRILADA. Call your HCP or get medical help right away if you have any of these symptoms of a serious allergic reaction:

- Hives
- Trouble breathing
- Swelling of your face, eyes, lips or mouth

Call your HCP or get medical care right away if you develop any of the above symptoms. Your treatment with ABRILADA may be stopped.

Please see Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including **BOXED WARNING**, Medication Guide, and Instructions for Use, available at ABRILADA.com.

Are there any side effects I should know about before starting treatment with ABRILADA? (continued)

ABRILADA can cause serious side effects, including: (continued)

• Nervous system problems

Signs and symptoms of a nervous system problem include: numbness or tingling, problems with your vision, weakness in your arms or legs, and dizziness.

• Blood problems

Your body may not make enough of the blood cells that help fight infections or help to stop bleeding. Symptoms include a fever that does not go away, bruising or bleeding very easily, or looking very pale.

• New heart failure or worsening of heart failure you already have. Call your healthcare professional right away if you get new or worsening symptoms of heart failure while taking ABRILADA, including:

- Shortness of breath
- Swelling of your ankles
- Sudden weight gain
- or feet

• Immune reactions including a lupus-like syndrome. Symptoms include chest discomfort or pain that does not go away, shortness of breath, joint pain, or a rash on your cheeks or arms that gets worse in the sun. Symptoms may improve when you stop ABRILADA

• Liver problems

Call your healthcare professional right away if you have any of these symptoms:

- Feel very tired
- Pain on the right side of your stomach (abdomen)
- Poor appetite or vomiting
- Skin or eyes look yellow

• Psoriasis

Some people using adalimumab products had new psoriasis or worsening of psoriasis they already had. Tell your HCP if you develop red scaly patches or raised bumps that are filled with pus. Your HCP may decide to stop your treatment with ABRILADA.

Call your HCP or get medical care right away if you develop any of the above symptoms. Your treatment with ABRILADA may be stopped.

The most common side effects of ABRILADA include:

- Injection site reactions: redness, rash, swelling, itching, or bruising. These symptoms usually will go away within a few days. Call your HCP right away if you have pain, redness or swelling around the injection site that does not go away within a few days or gets worse
- Upper respiratory infections (including sinus infections)
- Headaches
- Rash

These are not all of the possible side effects with ABRILADA. Tell your HCP if you have any side effect that bothers you or that does not go away. Ask your HCP or pharmacist for more information.



Not actual patient or healthcare professional.

What should I do if I experience side effects?

Please contact the doctor's office right away if you develop any symptoms of an infection. Also let your doctor know if you have any new or worsening medical conditions. Make sure to report any bruising, bleeding, persistent fever, or symptoms of infection. Seek immediate medical attention if you develop a severe allergic reaction.

You are encouraged to report adverse events related to Pfizer products by calling 1-800-438-1985 (US only). If you prefer, you may contact the US Food and Drug Administration (FDA) directly. Visit www.fda.gov/MedWatch or call 1-800-FDA-1088.

Please see the [Medication Guide](#) for more information.

Should I avoid any other medications while taking ABRILADA?

Tell your HCP about all of the medicines you take, including prescription medications, over-the-counter medicines, vitamins, and herbal supplements. Especially tell your HCP if you use:

- Orencia[®] (abatacept), Kineret[®] (anakinra), Remicade[®] (infliximab), Enbrel[®] (etanercept), Cimzia[®] (certolizumab pegol), or Simponi[®] (golimumab) because you should not use ABRILADA while you are also using one of these medicines
- Rituxan[®] (rituximab). Your HCP may not want to give you ABRILADA if you have recently received Rituxan
- Imuran[®] (azathioprine) or Purinethol[®] (6-mercaptopurine, 6-MP)

Please see Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including **BOXED WARNING**, Medication Guide, and Instructions for Use, available at ABRILADA.com.



Not actual patients.

SELECTED SAFETY INFORMATION

Common side effects of ABRILADA include injection site reactions (pain, redness, rash, swelling, itching, or bruising), **upper respiratory infections** (including sinus infections), **headaches**, and **rash**. These are not all of the possible side effects with ABRILADA. Tell your doctor if you have any side effect that bothers you or that does not go away.

Remember, tell your doctor right away if you have an infection or symptoms of an infection, including:

- Fever, sweats, or chills
- Muscle aches
- Cough
- Shortness of breath
- Blood in phlegm
- Weight loss
- Warm, red, or painful skin or sores on your body
- Diarrhea or stomach pain
- Burning when you urinate
- Urinating more often than normal
- Feeling very tired

ABRILADA is given by injection under the skin.

This is the most important information to know about ABRILADA. For more information, talk to your healthcare professional.

USES

ABRILADA is a prescription medicine used:

- **To reduce the signs and symptoms of:**
 - **Moderate to severe rheumatoid arthritis (RA) in adults.** ABRILADA can be used alone, with methotrexate, or with certain other medicines. ABRILADA may prevent further damage to your bones and joints and may help your ability to perform daily activities.
 - **Moderate to severe polyarticular juvenile idiopathic arthritis (JIA) in children** 2 years of age and older. ABRILADA can be used alone or with methotrexate.

- **Psoriatic arthritis (PsA) in adults.** ABRILADA can be used alone or with certain other medicines. ABRILADA may prevent further damage to your bones and joints and may help your ability to perform daily activities.
- **Ankylosing spondylitis (AS) in adults.**
- **Moderate to severe hidradenitis suppurativa (HS) in adults.**
- **To treat moderate to severe Crohn's disease (CD) in adults and children 6 years of age and older.**
- **To treat moderate to severe ulcerative colitis (UC) in adults.** It is not known if adalimumab products are effective in people who stopped responding to or could not tolerate anti-TNF medicines.
- **To treat moderate to severe chronic plaque psoriasis (Ps) in adults** who are ready for **systemic therapy or phototherapy, and are under the care of a doctor who will decide if other systemic therapies are less appropriate.**
- **To treat non-infectious intermediate** (middle part of the eye), **posterior** (back of the eye), and **panuveitis** (all parts of the eye) in adults.

Please see the Full Prescribing Information, including the Medication Guide, for ABRILADA.

You are encouraged to report adverse events related to Pfizer products by calling 1-800-438-1985 (U.S. only). If you prefer, you may contact the U.S. Food and Drug Administration (FDA) directly. Visit <http://fda.gov/MedWatch> or call 1-800-FDA-1088.

*Please see Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including **BOXED WARNING, Medication Guide, and Instructions for Use, available at [ABRILADA.com](http://Abrilada.com).***

INJECTION
Abrilada[™]
adalimumab-afzb



Not actual patient or healthcare professional.

Insurance coverage and financial support for your ABRILADA prescription

Is ABRILADA covered by my insurance?

ABRILADA may be covered by your insurance, which can be checked prior to your first injection.

Pfizer enCompass[®] offers a variety of support resources to help assess your insurance coverage and answer additional coverage-related questions you may have.



Pfizer enCompass provides access and reimbursement resources, support, and educational materials for ABRILADA.

A team of trained Access Counselors is available to help you understand your insurance benefits and to work with your HCP and health insurer on your behalf to access treatment with ABRILADA.

To assess your insurance coverage for ABRILADA or your support needs, **please contact an Access Counselor at 1-844-722-6672**, Monday–Friday, 8 AM to 8 PM ET, or visit www.PFIZERenCompass.com.

Please see Selected Safety Information throughout, Indications on page 3, and Full Prescribing Information, including BOXED WARNING, Medication Guide, and Instructions for Use, available at ABRILADA.com.

Pfizer enCompass Co-Pay Assistance Program for ABRILADA

The Pfizer enCompass Co-Pay Assistance Program for ABRILADA provides eligible, commercially insured individuals with a maximum benefit of \$4,000-\$14,000 per calendar year for out-of-pocket expenses.

If eligible, you may pay as little as \$0 for each ABRILADA treatment.

Federal and State health care beneficiaries not eligible. Private insurance only. The co-pay program covers only drug costs, not procedures, administration fees, or office visits. Enrollment required. Please see full Terms and Conditions* below.

***Terms and Conditions:** By using this program, you acknowledge that you currently meet the eligibility criteria and will comply with the terms and conditions below:

The Pfizer enCompass Co-Pay Assistance Program for ABRILADA is not valid for patients that are enrolled in a state or federally funded insurance program, including but not limited to Medicare, Medicaid, TRICARE, Veterans Affairs health care, a state prescription drug assistance program, or the Government Health Insurance Plan available in Puerto Rico (formerly known as “La Reforma de Salud”). Program offer is not valid for cash-paying patients. Patients prescribed ABRILADA for adolescent hidradenitis suppurativa (HS), pediatric uveitis, or pediatric ulcerative colitis are not eligible for this co-pay savings program. With this program, eligible patients may pay as little as \$0 co-pay per ABRILADA treatment, subject to a maximum benefit of \$4,000-\$14,000 per calendar year for out-of-pocket expenses for ABRILADA, depending on your insurance, including co-pays or coinsurances. The amount of any benefit is the difference between your co-pay and \$0. After the maximum benefit, you will be responsible for the remaining monthly out-of-pocket costs. Patient must have private insurance with coverage of ABRILADA. This offer is not valid when the entire cost of your prescription drug is eligible to be reimbursed by your private insurance plans or other private health or pharmacy benefit programs. You must deduct the value of this assistance from any reimbursement request submitted to your private insurance plan, either directly by you or on your behalf. You are responsible for reporting use of the program to any private insurer, health plan, or other third party who pays for or reimburses any part of the prescription filled using the program, as may be required. You should not use the program if your insurer or health plan prohibits use of manufacturer co-pay assistance programs. This program is not valid where prohibited by law. This program cannot be combined with any other savings, free trial or similar offer for the specified prescription. **Co-pay card will be accepted only at participating pharmacies. If your pharmacy does not participate, you may be able to submit a request for a rebate in connection with this offer. The rebate form can be found at <https://patient.pfizerandicopay.com>. This program is not health insurance.** This program is good only in the U.S. and Puerto Rico. This program is limited to 1 per person during this offering period and is not transferable. This offer cannot be redeemed more than once per 30 days per patient. No other purchase is necessary. Data related to your redemption of the program assistance may be collected, analyzed, and shared with Pfizer, for market research and other purposes related to assessing Pfizer’s programs. Data shared with Pfizer will be aggregated and de-identified; it will be combined with data related to other assistance redemptions and will not identify you. Pfizer reserves the right to rescind, revoke or amend this program without notice. This program may not be available to patients in all states. For more information about Pfizer, visit www.pfizer.com. For more information about the Pfizer enCompass Co-Pay Assistance Program, call Pfizer enCompass at 1-844-722-6672, or write to Pfizer enCompass Co-Pay Assistance Program 2730 S. Edmonds Lane, Suite 300, Lewisville, TX. Card and Program expires 12/31/2024.



Is there a mobile app that can help me manage my therapy?

The **All in 1** app is a free mobile app—designed to help you keep track of your injections and symptoms along your treatment journey.

All in 1

The **All in 1** app includes:

- A **customizable calendar** allowing you to keep track of injection details and receive important reminders
- 1-on-1 education to help understand proper injection techniques from **Nurse Guides***
- A simple way to **keep track of key details**—such as your symptoms—that you can share with your doctor

To download the **All in 1** app, visit your mobile app store.



App Store is a service mark of Apple Inc., registered in the U.S. and other countries. Google Play and the Google Play logo are trademarks of Google LLC.

*Enrollment in Pfizer enCompass[®] required.

To learn more about ABRILADA,
visit www.ABRILADA.com

The brand names included throughout are the properties of their respective owners.